



Comparison of metoprolol degradation by Fe^{III}-NTA modified Fenton-like reaction in the absence and presence of manganese: Efficiency and intermediates




Yifan Li^a, Jianhui Sun^a, Sheng-Peng Sun^{b,*}

^a K L E, w R H R E P C, M E, H K L E P C
^b E, H N, H 453007, P R C C E E, C C, C E M
 w, J 215123, P R C

HIGHLIGHTS

- Metoprolol degradation by Fe^{III}-NTA/H₂O₂ was enhanced in the presence of Mn^{II}.
- Rapid degradation of metoprolol was obtained by Fe^{III}-NTA/H₂O₂-Mn at pH 4.0–8.0.
- Fe^{III}-NTA/H₂O₂-Mn led to more efficient degradation of metoprolol and intermediates.
- Degradation pathways of metoprolol were proposed.

GRAPHICAL ABSTRACT



ARTICLE INFO

A :
 Received 7 October 2016
 Received in revised form 21 December 2016
 Accepted 22 December 2016
 Available online 26 December 2016

K w :
 Metoprolol
 Fe^{III}-NTA
 Fenton-like reaction
 Hydroxyl radical
 Manganese

ABSTRACT

The degradation of metoprolol in aqueous solutions by traditional Fenton-like reaction (Fe^{III}/H₂O₂), and Fe^{III}-NTA modified Fenton-like reaction (Fe^{III}-NTA/H₂O₂) in the absence and presence of Mn^{II} (Fe^{III}-NTA/H₂O₂-Mn) have been investigated. The results show that Fe^{III}-NTA/H₂O₂ is able to degrade metoprolol at initial neutral pH. In particular, the presence of Mn^{II} greatly improved the degradation rate of metoprolol by Fe^{III}-NTA/H₂O₂ over a wide pH range of 4.0–8.0. Under the same conditions, the degradation rate constants of metoprolol (k) obtained in Fe^{III}-NTA/H₂O₂-Mn system were typically 7–9-fold larger than those obtained in Fe^{III}-NTA/H₂O₂ system. The involved reactions in Fe^{III}-NTA/H₂O₂-Mn system were proposed on the basis of important parameters analysis including ferrous ion concentration, dissolved oxygen (DO) concentration, and the quenching experiments for hydroxyl radical (HO·) and superoxide anion radical (O₂⁻). Several intermediates have been identified by mass spectrometry. Our results suggest that the degradation of metoprolol in both Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems were caused by HO· attack. The degradation pathways of metoprolol were proposed on the basis of the identified intermediates. Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Metoprolol, a selective β₁ receptor blocking drug, is widely used in the therapy of hypertension and angina. In the past decade, metoprolol and its metabolites have been detected in sewage treatment plant effluents, surface water, groundwater, and even

* Corresponding author.

E-mail address: shepsun@suda.edu.cn (S.-P. Sun).

in drinking water, which pose a potential risk to human health [1,2]. The degradation rate of metoprolol in water by the direct photolysis is small, e.g., the extrapolated half-lives of metoprolol in central Europe and the U.S. were reported at 28–95 d in summer and 190–449 d in winter [3]. Thus, metoprolol is most likely to be pseudo-persistent in natural waters if do not consider the indirect photolysis effects. In addition, the efficiency of metoprolol removal is lower than 30% in traditional sewage treatment plant [4]. There is an urgent need to improve the removal efficiencies of pharmaceuticals in water.

A far more effective method is the so-called advanced oxidation processes (AOPs), involving the generation of hydroxyl radical ($\text{HO}\cdot$) [5–7], have been widely applied in water treatment for harmful organic compounds removal. Efficient degradation of metoprolol has been obtained by photocatalysis, UV/ H_2O_2 and Fenton-related processes [8–10]. Two decades ago, research by Sun and Pignatello [11] showed the successful degradation of 2,4-dichlorophen-oxycetic acid at pH 6.0 by chelates modified Fenton-like reaction. Chelating agents modified Fenton-like reaction has been receiving increasing attention because it can work at neutral pH conditions, which overcomes the limitation of acidic pH (e.g., optimal pH 2.8–3.0) by using traditional Fenton and/or Fenton-like reaction. S,S-ethylenediamine-N,N'-disuccinic acid (EDDS) modified Fenton reaction (i.e., $\text{Fe}^{\text{II}}\text{-EDDS}/\text{H}_2\text{O}_2$) and Fenton-like reaction (i.e., $\text{Fe}^{\text{III}}\text{-EDDS}/\text{H}_2\text{O}_2$) are capable of degrading 4-chlorophenol, bisphenol A, naphthenic acid, and 4-tert-butylphenol at near neutral or basic pH conditions [12–15]. The use of $\text{Fe}^{\text{III}}\text{-EDDS}$ for photo-Fenton reaction at neutral pH is also successful in degrading micro-pollutants in wastewater treatment plant effluents [16–18]. In addition, nitrilotriacetic acid (NTA) and ethylenediaminetetraacetic acid (EDTA) modified Fenton-like reactions (i.e., $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-EDTA}/\text{H}_2\text{O}_2$) have also been proved to be highly effective in degrading atrazine, carbamazepine, fenuron, naphthenic acid, parachlorobenzoic acid and sulfamethoxazole at neutral or slightly basic pH conditions [19–22]. In particular, our recent finding suggested that the presence of manganese (Mn^{II}) could greatly enhance the Fenton-like catalytic activity of $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ [23]. The potential mechanism may involve an enhanced generation of superoxide anion radical ($\text{O}_2^{\cdot-}$) by the Mn^{II} , which can accelerate the reduction of $\text{Fe}^{\text{III}}\text{-NTA}$ to $\text{Fe}^{\text{II}}\text{-NTA}$ and indirectly improve the generation of $\text{HO}\cdot$ [23]. The $\text{Mn}^{\text{II}}/\text{Fe}^{\text{III}}\text{-NTA}$ catalytic system would be very promising since Fe^{III} and Mn^{II} are quite ubiquitous in the environment and can be used as natural Fenton-like catalyst in the presence of NTA (or natural Fe^{III} complexing agents) and H_2O_2 . However, little research on this system has been reported.

The purpose of this study is to demonstrate the potential effectiveness of $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ for water treatment. In the present study, we choose metoprolol as a target compound, to evaluate the efficiency and intermediates generated from metoprolol degradation by $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ in the absence and presence of Mn^{II} . The degradation kinetics of metoprolol in both systems were compared under various conditions. The involved reactions in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system were proposed on the basis of certain important parameters analysis such as ferrous ion concentration, dissolved oxygen (DO) concentration, and the quenching experiments for $\text{HO}\cdot$ and $\text{O}_2^{\cdot-}$. The intermediates were identified by liquid chromatography/mass spectrometry.

2. Materials and methods

2.1. Chemicals

Ferric chloride hexahydrate, hydrogen peroxide (30% H_2O_2 , w/w), sodium hydroxide and sulfuric acid were purchased from Sino-

pharm Chemical Reagent Co., Ltd. Metoprolol tartrate (purity $\geq 98\%$) was purchased from LKT Laboratories, Inc. NTA was purchased from Acros Organics. Formic acid and manganese nitrate tetrahydrate were purchased from Sigma-Aldrich. HPLC-grade methanol, 2-propanol and water were purchased from Fisher Scientific. Chloroform was purchased from Chinasun Specialty Products Co., Ltd. All chemicals were of reagent grade or better, and used as received without further purification. Milli-Q water was used to prepare solutions.

2.2. Experimental procedure

All experiments were performed in 125 ml glass Erlenmeyer flasks. Briefly, 100 ml of metoprolol aqueous solution and an appropriate amount of freshly prepared $\text{Fe}^{\text{III}}\text{-NTA}$ solution were added into the flask. The flask was then placed in a water-jacketed glass vessel water bath ($25 \pm 1^\circ\text{C}$) under a magnetic stirring condition. After that, an appropriate amount of Mn^{II} was added to the solution. The pH of the solution was then adjusted to the desired value using 1.0/0.1 M NaOH or 1.0/0.1 M H_2SO_4 . Finally, Fenton-like reaction was started by the addition of an appropriate amount of H_2O_2 . It is worth noting that the water-jacketed glass vessel was covered with aluminum foil to prevent the effect of light on the Fenton-like reaction. Samples (1 ml) were withdrawn at different time intervals and mixed with 0.1 ml of methanol immediately to quench the reaction.

2.3. Instrumentation

The concentration of metoprolol was determined by a high performance liquid chromatography (HPLC, Agilent 1100) with fluorescence detection. An Ultra C-18 column ($5 \mu\text{m}$, $250 \times 4.6 \text{ mm}$) was used for the chromatographic separation. The mobile phase consisted of methanol and 0.1% formic acid aqueous solution (35:65, v/v) at a flow rate of 0.9 ml min^{-1} . The column temperature was 30°C . The injection volume was $20 \mu\text{l}$, and the fluorescence detection was performed at 275 nm of excitation and 298 nm of emission.

The degradation intermediates were identified by an Agilent 1200 HPLC system coupled to a 6130 quadrupole mass spectrometer. The MS detector was performed for a full scan range of 30–600 under an ESI + APCI positive mode. The MS operating conditions were as follows: Fragmentor 90, drying gas flow rate 6 L min^{-1} , nebulizer pressure 45 psi, dry gas temperature 300°C , vaporizer temperature 250°C , capillary voltage 3500 V, corona current $4.0 \mu\text{A}$ and charging voltage 2000 V.

The concentration of ferrous ion in the aqueous solutions was determined by using the 1,10-phenanthroline method. The concentration of H_2O_2 was determined by the iodide method [24]. The dissolved oxygen (DO) concentration in the aqueous solutions during the reaction time was monitored by a dO_2 sensor (Model Z010023525, AppliSens).

3. Results and discussion




3.1. Comparison of metoprolol degradation by $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems

Fig. 1 shows a comparison of metoprolol degradation by $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems at initial neutral pH. A minimum degradation of metoprolol was observed in $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$ system, with 3.61% degradation efficiency of metoprolol after 60 min of the reaction time. This result was expected because of the precipitation of ferric ion at pH 7.0 in the absence of chelator. In contrast, $\text{Fe}^{\text{III}}\text{-NTA}$ complex is able to catalyze

Fenton-like reaction at pH 7.0, resulted in 100% degradation efficiency of metoprolol after 30 min of the reaction time in Fe^{III}-NTA/H₂O₂ system. In particular, Fe^{III}-NTA/H₂O₂-Mn system led to a very fast degradation of metoprolol, with 100% degradation efficiency of metoprolol after 7 min of the reaction time. The result is consistent with our recent finding that the presence of Mn^{II} could enhance Fe^{III}-NTA-catalyzed Fenton-like reaction [23].

The effects of Mn^{II} concentration on the degradation kinetics of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were shown in Fig. 2. An increase in Mn^{II} concentration from 0.01 to 0.1 mM, corresponding to Mn:Fe molar ratios of 0.1:1 to 1:1, had a positive effect on metoprolol degradation. The apparent degradation rate constant of metoprolol (k_{app}) increased from 0.7547 to 1.6048 min⁻¹. However, further increasing Mn^{II} concentration from 0.2 to 1.0 mM led to a slight inhibitory effect on the degradation rate of metoprolol, e.g., the k_{app} value of metoprolol decreased from 1.6289 to 1.1103 min⁻¹. This is because when Mn^{II} concentration was larger than 0.2 mM, hydrous manganese dioxide could form from free Mn^{II} in excess

from the reaction of Fe^{III}-NTA with H₂O₂ in Fe^{III}-NTA/H₂O₂ system (i.e., R7), thus the generation rate of Fe^{II}-NTA in Fe^{III}-NTA/H₂O₂ system is much slower than that in Fe^{III}-NTA/H₂O₂-Mn system.



the range of NTA:Fe molar ratios tested, the values of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were 7–17-fold larger than those in Fe^{III}-NTA/H₂O₂ system. An increase in NTA/Fe molar ratio from 0.5:1 to 2:1 had positive effects on the degradation rates of metoprolol in both Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems. The values of metoprolol increased from 0.0311 to 0.1754 min⁻¹ in Fe^{III}-NTA/H₂O₂ system and increased from 0.3450 to 1.6048 min⁻¹ in Fe^{III}-NTA/H₂O₂-Mn system, respectively. When NTA/Fe molar ratio was larger than 2:1, however, a slight decrease in the values of metoprolol was observed in either Fe^{III}-NTA/H₂O₂ or Fe^{III}-NTA/H₂O₂-Mn system. This can be explained by the fact that the free NTA in excess scavenges HO·. Although the reaction rate constant of metoprolol with HO· (5.2–8.4 × 10⁹ M⁻¹ s⁻¹) [32,33] is larger than that of Fe^{III}-NTA/free NTA with HO·, it is believed that the competitive consumption of HO· by the free NTA in excess can slightly limit the degradation of metoprolol.

3.4. E H₂O₂:F^{III}

Fig. 6 shows the effects of H₂O₂/Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems at initial neutral pH. Without the addition of H₂O₂, no degradation of metoprolol was observed in both systems. Over the range of H₂O₂/Fe^{III} molar ratios tested, the values of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were 5–9-fold larger than those in Fe^{III}-NTA/H₂O₂ system. With increasing H₂O₂/Fe^{III} molar ratio from 10:1 to 200:1, the values of metoprolol continuously increased from 0.0598 to 0.2276 min⁻¹ in Fe^{III}-NTA/H₂O₂ system. In Fe^{III}-NTA/H₂O₂-Mn system, the values of metoprolol increased greatly from 0.3384 to 1.6048 min⁻¹ and reached a plateau with increasing H₂O₂/Fe^{III} molar ratio from 10:1 to 100:1. It is expected that the use of higher molar ratio of H₂O₂/Fe^{III} led to higher rate of HO· generation, by which faster degradation of metoprolol could be obtained. No inhibitory effect of H₂O₂ on the degradation kinetics of metoprolol was observed even at H₂O₂/Fe^{III} molar ratio of 200:1. This is because the reaction rate constant of H₂O₂ with HO· ((1.2–4.5) × 10⁷ M⁻¹ s⁻¹) [34] is much smaller than those of metoprolol, Fe^{III}-NTA and free NTA with HO·.

3.5. E F^{III}

Fe^{III} acts as the catalyst to initiate Fenton-like reaction, a high rate of HO· generation might be obtained at high concentration of Fe^{III}-NTA. The effects of Fe^{III} dosages on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems at




Fig. 6. Effects of NTA/Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III}, NTA/Fe^{III} molar ratio of 0.5:1–5:1, H₂O₂/Fe^{III} molar ratio of 100:1, Mn^{II}/Fe^{III} molar ratio of 1:1, and initial pH of 7.0.




Fig. 7. Effects of H₂O₂/Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III}, NTA/Fe^{III} molar ratio of 2:1, H₂O₂/Fe^{III} molar ratio of 5:1–200:1, Mn^{II}/Fe^{III} molar ratio of 1:1, and initial pH of 7.0.




Fig. 8. Effects of Fe^{III} dosages on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.01–0.15 mM of Fe^{III}, NTA/Fe^{III} molar ratio of 2:1, H₂O₂/Fe^{III} molar ratio of 100:1, Mn^{II}/Fe^{III} molar ratio of 1:1, and initial pH of 7.0.




Fig. 9. Effects of H₂O₂/Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems at initial neutral pH were shown in Fig. 7. No degradation of metoprolol was observed in the absence of Fe^{III}, which suggests that HO· is generated from the Fenton-like reaction of Fe^{III}/NTA/H₂O₂. Over

the range of Fe^{III} concentrations tested, the k values of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were 7–9-fold larger than that in Fe^{III}-NTA/H₂O₂ system. An increase in Fe^{III} concentrations from 0 to 0.15 mM has positive effects on the degradation rate of metoprolol in both systems. The k values of metoprolol increased from

0 to 0.3174 min⁻¹ in Fe^{III}-NTA/H₂O₂ system and to 2.2988 min⁻¹ in Fe^{III}-NTA/H₂O₂-Mn system, respectively. The results indicate that the rate of HO[•]

Fe^{III}-NTA would increase the generation rate of Fe^{II}-NTA (R5) and (R7), which in turn reacts with H₂O₂ to produce more HO[•].

3.6. I

w

It was worth noting that a high initial concentration of metoprolol (292 μM, ten times higher than that in typical experiments) was used in order to capture more intermediates. The typical full-scan total molecular ion chromatograms of samples at different reaction time (RT) can be found in the [Supplementary data](#). Two major peaks with molecular ion ([M+H]⁺) at mass-to-charge (/) ratios of 268 and 192 were observed in samples before the treatment, corresponding to metoprolol and NTA, respectively. Several intermediates were detected in samples after the reaction by mass spectrometry, their molecular ion were identified at / 134, 238, 258, 274, 282, 284, 298, 300, 314, 316, 330, 332 and 348. A major degradation product is 3-(isopropylamino)propane-1,2-diol (/ 134), which formed by the cleavage of aromatic ether bond of metoprolol due to the attack of HO[•]. A same intermediate has been detected in the degradation of metoprolol by photocatalysis, γ-irradiation, Fenton and photo-Fenton processes [8,10,33,35]. In addition, the attack of HO[•] on the ether bond and/or the β-C in methoxyethyl side chain of metoprolol gave intermediates with molecular ion at / 238, corresponding to 4-(2-hydroxy-3-(isopropylamino)propoxy)benzaldehyde and 1-(4-ethylphenoxy)-3-(isopropylamino)propan-2-ol, consistent with previous studies [8,10].

The hydroxylation of benzene ring (and/or side chain) of metoprolol led to the formation of hydroxyl-metoprolol isomers (/ 284). The intermediates with molecular ion at / 282 could be 4-(2-methoxyethyl)phenyl 2-hydroxy-3-(isopropylamino)propanoate and 1-(4-(2-hydroxy-3-(isopropylamino)propoxy)phenyl)-2-methoxyethanone, which were formed by the oxidation of hydroxyl group to keto group in hydroxyl-metoprolol (i.e., 3-(isopropylamino)-1-(4-(2-methoxyethyl)phenoxy)propane-1,2-diol and 1-(4-(1-hydroxy-2-methoxyethyl)phenoxy)-3-(isopropylamino)propan-2-ol). Moreover, the addition of 1 HO[•], 2 HO[•], 3 HO[•] and 4 HO[•] on hydroxyl-metoprolol gave intermediates with molecular ion at / 300, 316, 332 and 348, corresponding to di-, tri-, tetra-, and penta-hydroxyl-metoprolol, respectively. The intermediates with molecular ion at / 298 and 314 could be formed by the addition of 1 HO[•] and 2 HO[•] on the detected intermediates (/ 282). The intermediates with molecular ion at / 258 and 274 were probably formed by the addition of 2 HO[•] and 3 HO[•] on two undetected intermediates (/ 226), i.e., 1-amino-3-(4-(2-methoxyethyl)phenoxy)propan-2-ol and 4-(2-hydroxy-3-(isopropylamino)propoxy)phenol [33,36].

A comparison of Figs. S2 and S3 demonstrates that Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates. In addition, Fig. 8 shows the changes in the total molecular ion chromatogram area of metoprolol and its intermediates with the reaction time. It can be seen that the total areas of major intermediates (i.e., / 238, 282, 284, 298, 300, etc.) increased rapidly to reach their maximum values within the first 3–5 min of the reaction time, and then decreased to undetectable levels at 20 min of the reaction time. In particular, the major intermediate (/ 134) increased continuously until almost complete degradation of metoprolol took place (12–15 min of the reaction time), and then began to decrease slowly over time. The degradation pathways of metoprolol were proposed on the basis of these identified intermediates, which were presented in Scheme 1. Our results clearly suggested that the degradation of metoprolol and its intermediates were mainly caused by HO[•] attack.

4.

The results of this study indicate that Fe^{III}-NTA/H₂O₂ is able to degrade metoprolol at initial neutral pH, and the degradation rate

of metoprolol can be greatly improved with the presence of Mn^{II}. The values of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were typically 7–9-fold larger than those in Fe^{III}-NTA/H₂O₂ system. It can be concluded that the presence of Mn^{II} does not directly catalyze Fenton-like reaction, but indirectly enhance Fe^{III}-NTA-catalyzed Fenton-like reaction over a wide pH range of 4.0–8.0. The main intermediates during the degradation of metoprolol have been identified by mass spectrometry, which suggested that the degradation of metoprolol and its intermediates were mainly caused by HO[•] attack. Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates.

This work was supported in part by the Innovation Scientists and Technicians Troop Construction Projects of Henan Province, Plan for Scientific Innovation Talent of Henan Province (Grant No. 134200510014), and in part by the Natural Science Foundation of Jiangsu Province (Grant No. BK20140341). The authors also would like to thank anonymous reviewers for their thoughtful comments and suggestions on this manuscript.

Quenching tests with 2-propanol and chloroform on the degradation of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system (Fig. S1). Total molecular ion chromatogram of metoprolol and its intermediates in Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems (Fig. S2 and Fig. S3). Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.cej.2016.12.098>.

- [1] M.S. Kostich, A.L. Batt, J.M. Lazorchak, Concentrations of prioritized pharmaceuticals in effluents from 50 large wastewater treatment plants in the US and implications for risk estimation, *Environ. Pollut.* 184 (2014) 354–359.
- [2] J. Sun, Q. Luo, D. Wang, Z. Wang, Occurrences of pharmaceuticals in drinking water sources of major river watersheds, China, *Ecotoxicol. Environ. Saf.* 117 (2015) 132–140.
- [3] Q.-T. Liu, H.E. Williams, Kinetics and degradation products for direct photolysis of β-blockers in water, *Environ. Sci. Technol.* 41 (2007) 803–810.
- [4] C. Miège, J.M. Choubert, L. Ribeiro, M. Eusèbe, M. Coquery, Fate of pharmaceuticals and personal care products in wastewater treatment plants – conception of a database and first results, *Environ. Pollut.* 157 (2009) 1721–1726.
- [5] C.-C. Chen, R.-J. Wu, Y.-Y. Tzeng, C.-S. Lu, Chemical oxidative degradation of acridine orange dye in aqueous solution by Fenton's reagent, *J. Chin. Chem. Soc.* 56 (2009) 1147–1155.
- [6] H.-J. Fan, S.-T. Huang, W.-H. Chung, J.-L. Jan, W.-Y. Lin, C.-C. Chen, Degradation pathways of crystal violet by Fenton and Fenton-like systems: condition optimization and intermediate separation and identification, *J. Hazard. Mater.* 171 (2009) 1032–1044.
- [7] C.-C. Chen, W.-C. Chen, M.-R. Chiou, S.-W. Chen, Y.Y. Chen, H.-J. Fan, Degradation of crystal violet by an FeGAC/H₂O₂ process, *J. Hazard. Mater.* 196 (2011) 420–425.
- [8] B. Abramović, S. Kler, D. Šojić, M. Laušević, T. Radović, D. Vione, Photocatalytic degradation of metoprolol tartrate in suspensions of two TiO₂-based photocatalysts with different surface area. Identification of intermediates and proposal of degradation pathways, *J. Hazard. Mater.* 198 (2011) 123–132.
- [9] F.J. Rivas, O. Gimeno, T. Borralho, M. Carbajo, UV-C radiation based methods for aqueous metoprolol elimination, *J. Hazard. Mater.* 179 (2010) 357–362.
- [10] V. Romero, S. Acevedo, P. Marco, J. Giménez, S. Esplugas, Enhancement of Fenton and photo-Fenton processes at initial circumneutral pH for the degradation of the β-blocker metoprolol, *Water Res.* 88 (2016) 449–457.
- [11] Y. Sun, J.J. Pignatello, Chemical treatment of pesticide wastes. Evaluation of iron(III) chelates for catalytic hydrogen peroxide oxidation of 2,4-D at circumneutral pH, *J. Agric. Food Chem.* 40 (1992) 322–327.
- [12] A. Rastogi, S.R. Al-Abed, D.D. Dionysiou, Effect of inorganic, synthetic and naturally occurring chelating agents on Fe(II) mediated advanced oxidation of chlorophenols, *Water Res.* 43 (2009) 684–694.
- [13] W. Huang, M. Brigante, F. Wu, C. Mousty, K. Hanna, G. Mailhot, Assessment of the Fe(III)-EDDS complex in Fenton-like processes: from the radical formation to the degradation of bisphenol A, *Environ. Sci. Technol.* 47 (2013) 1952–1959.

- [14] Y. Wu, M. Passananti, M. Brigante, W. Dong, G. Mailhot, Fe(III)-EDDS complex in Fenton and photo-Fenton processes: from the radical formation to the degradation of a target compound, *Environ. Sci. Pollut. Res.* 21 (2014) 12154–12162.
- [15] Y. Zhang, N. Klammerth, S.A. Messele, P. Chelme-Ayala, M. Gamal El-Din, Kinetics study on the degradation of a model naphthenic acid by ethylenediamine-N, N'-disuccinic acid-modified Fenton process, *J. Hazard. Mater.* 318 (2016) 371–378.
- [16] N. Klammerth, S. Malato, A. Agüera, A. Fernández-Alba, G. Mailhot, Treatment of municipal wastewater treatment plant effluents with modified photo-Fenton as a tertiary treatment for the degradation of micro pollutants and disinfection, *Environ. Sci. Technol.* 46 (2012) 2885–2892.
- [17] N. Klammerth, S. Malato, A. Agüera, A. Fernández-Alba, Photo-Fenton and modified photo-Fenton at neutral pH for the treatment of emerging contaminants in wastewater treatment plant effluents: a comparison, *Water Res.* 47 (2013) 833–840.
- [18] S. Miralles-Cuevas, I. Oller, J.A.S. Pérez, S. Malato, Removal of pharmaceuticals from MWTP effluent by nanofiltration and solar photo-Fenton using two different iron complexes at neutral pH, *Water Res.* 64 (2014) 23–31.
- [19] Y.H. Dao, J. De Laat, Hydroxyl radical involvement in the decomposition of hydrogen peroxide by ferrous and ferric-nitritotriacetate complexes at neutral pH, *Water Res.* 45 (2011) 3309–3317.
- [20] S.-P. Sun, X. Zeng, A.T. Lemley, Kinetics and mechanism of carbamazepine degradation by a modified Fenton-like reaction with ferric-nitritotriacetate complexes, *J. Hazard. Mater.* 252–253 (2013) 155–165.
- [21] Y. Zhang, N. Klammerth, M. Gamal El-Din, Degradation of a model naphthenic acid by nitritotriacetic acid – modified Fenton process, *Chem. Eng. J.* 292 (2016) 340–347.
- [22] A. De Luca, R.F. Dantas, S. Esplugas, Assessment of iron chelates efficiency for photo-Fenton at neutral pH, *Water Res.* 61 (2014) 232–242.
- [23] Y. Li, J. Sun, S.-P. Sun, Mn²⁺-mediated homogeneous Fenton-like reaction of Fe(III)-NTA complex for efficient degradation of organic contaminants under neutral conditions, *J. Hazard. Mater.* 313 (2016) 193–200.
- [24] C. Kormann, D.W. Bahnemann, M.R. Hoffmann, Photocatalytic production of hydrogen peroxides and organic peroxides in aqueous suspensions of titanium dioxide, zinc oxide, and desert sand, *Environ. Sci. Technol.* 22 (1988) 798–806.
- [25] R.J. Watts, J. Sarasa, F.J. Loge, A.L. Teel, Oxidative and reductive pathways in manganese-catalyzed Fenton's reactions, *J. Environ. Eng. ASCE* 131 (2005) 158–164.
- [26] B.H.J. Bielski, D.E. Cabelli, R.L. Arudi, A.B. Ross, Reactivity of HO₂/O₂⁻ radicals in aqueous solution, *J. Phys. Chem. Ref. Data* 14 (1985) 1041–1100.
- [27] J. De Laat, Y.H. Dao, N. Hamdi El Najjar, C. Daou, Effect of some parameters on the rate of the catalysed decomposition of hydrogen peroxide by iron(III)-nitritotriacetate in water, *Water Res.* 45 (2011) 5654–5664.
- [28] G.V. Buxton, C.L. Greenstock, W.P. Helman, A.B. Ross, Critical-review of rate constants for reactions of hydrated electrons, hydrogen-atoms and hydroxyl radicals (·OH/O⁻) in aqueous-solution, *J. Phys. Chem. Ref. Data* 17 (1988) 513–886.
- [29] R.J. Motekaitis, A.E. Martell, The iron(III) and iron(II) complexes of nitritotriacetic acid, *J. Coord. Chem.* 31 (1994) 67–78.
- [30] B.A. Smith, A.L. Teel, R.J. Watts, Identification of the reactive oxygen species responsible for carbon tetrachloride degradation in modified Fenton's systems, *Environ. Sci. Technol.* 38 (2004) 5465–5469.
- [31] M. Bucheli-Witschel, T. Egli, Environmental fate and microbial degradation of aminopolycarboxylic acids, *FEMS Microbiol. Rev.* 25 (2001) 69–106.
- [32] C. Slegers, G. Baldacchino, D. Le Parc, B. Hickel, B. Tilquin, Radical mechanisms in the radiosterilization of metoprolol tartrate solutions, *Pharm. Res.* 20 (2003) 1977–1983.
- [33] W. Song, W.J. Cooper, S.P. Mezyk, J. Greaves, B.M. Peake, Free radical destruction of β-blockers in aqueous solution, *Environ. Sci. Technol.* 42 (2008) 1256–1261.
- [34] C. Walling, Fenton's reagent revisited, *Acc. Chem. Res.* 8 (1975) 125–131.
- [35] H. Yang, T. An, G. Li, W. Song, W.J. Cooper, H. Luo, X. Guo, Photocatalytic degradation kinetics and mechanism of environmental pharmaceuticals in aqueous suspension of TiO₂: a case of β-blockers, *J. Hazard. Mater.* 179 (2010) 834–839.
- [36] J. Radjenovic, B.I. Escher, K. Rabaey, Electrochemical degradation of the β-blocker metoprolol by Ti/Ru_{0.7}Ir_{0.3}O₂ and Ti/SnO₂-Sb electrodes, *Water Res.* 45 (2011) 3205–3214.